8200076

AHIE UNIVIED SHAMES OF AMIERIOA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Frank Holland

Telhereas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART Hereof, and the various requirements of ${\rm LAW}$ in such cases made and provided have BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, therefore, this certificate of plant variety protection is to grant UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLI-CANT(S) FOR THE TERM OF eighteen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-CLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT RIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. HE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS SS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS BY THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

POPCORN

'Big Red'

In Testimony Withereof, I have hereunto set my hand and caused the seal of the Want Tariety Protection Office to be affixed at the City of washington, D.C.

this 30th day of December in the year of our Lord one thousand nine hundred and eighty-two

Plant Variety Protection Office Grain Division Agricultural Marketing Service

-	UNITED STATES DEPARTME AGRICULTURAL MARK LIVESTOCK, POULTRY, GRA	CETING SERVICE			FORM AP OMB NO.		
AF	PPLICATION FOR PLANT VARIESTRUCTIONS: See Reverse.		No certificate for plant variety protection may be issued unless a completed application form has been received (5 U.S.C. 553).				
	. TEMPORARY DESIGNATION OF VARIETY	1b, VARIETY NAME			CIAL USE ONLY		
	(FH-1)	BIG RED		PV NUMBER	2080		
2.	KIND NAME	3. GENUS AND SPE	CIES NAME	FILING DATE	10076	A.M.	
	Popcorn	Zea May	S	3/4/82	3:00	P.M.	
4.	FAMILY NAME (BOTANICAL)	5. DATE OF DETER		FEE RECEIVED	DATE		
		Sept. 19		\$ 500.00 \$ 250.00	3/4/82 12/10/82	?	
6.	NAME OF APPLICANT(S)	7. ADDRESS (Street	and No. or R.F.D. No.,	City, State, and ZIP	8. TELEPHON		
	Frank Holland		Old Logan Rer, Ohio 43130		614 65		
9.	IF THE NAMED APPLICANT IS NOT A PE	RSON, FORM OF	10. IF INCORPORATE		0455 11. DATE OF 1	NCOR-	
	ORGANIZATION: (Corporation, partnershi	ip, association, etc.)	DATE OF INCORE	PORATION	PORATION	1	
12.	NAME AND MAILING ADDRESS OF APPL	ICANT REPRESENTA	TIVE(S), IF ANY, TO S	ERVE IN THIS APPLI	CATION AND RE	CEIVE	
	ALL PAPERS:			ş		01 , 1	
٠.							
13.	CHECK BOX BELOW FOR EACH ATTACH	MENT SUBMITTED:				•	
8.A	13A. Exhibit A, Origin and Bree	ding History of the	Variety (See Section 5	2 of the Plant Varies	ty Protection Ac	t.)	
	X 13B. Exhibit B, Novelty Stateme		***			'/	
		72.5				-	
				Plant Variety Protec	tion Office.)		
	130 Exhibit D, Additional Desc	ription of the Variet	у.				
14a.	DOES THE APPLICANT(S) SPECIFY THAT	SEED OF THIS VARI	ETY BE SOLD BY VAR	IETY NAME ONLY A	S A CLASS OF CE	RTIFIED	
14b.	SEED? (See Section 83(a). (If "Yes," answe	•		NO			
140.	DOES THE APPLICANT(S) SPECIFY THAT LIMITED AS TO NUMBER OF GENERATION	THIS VARIETY BE DNS?	14c. IF "YES," TO 14E TION BEYOND B	3, HOW MANY GENEF REEDER SEED?	RATIONS OF PRO	DUC-	
	X YES NO		FOUNDATION	REGISTERED	CERTIFIEI)	
15a.	DID THE APPLICANT(S) FILE FOR PROTE name of countries and dates,)	CTION OF THIS VAR	IETY IN OTHER COUN	ITRIES? TYES	NO (If "Y	es," give	
					•		
15Ь.	HAVE RIGHTS BEEN GRANTED THIS VA and dates.)	RIETY IN OTHER CO	UNTRIES? YES	X NO (If "Yes,"	give name of cour	itries	
	and the second s	and the state of the state of	A Land to the second	e grande			
16.	DOES THE APPLICANT(S) AGREE TO THE JOURNAL?			***			
17.	The applicant(s) declare(s) that a viable replenished upon request in accordance	sample of basic seed with such regulation	of this variety will be s as may be applicable	furnished with the	application and	will be	
4.	The undersigned applicant(s) is (are) the owner(s) of this sexually reproduced novel plant variety, and believe(s) that the variety is distinct, uniform, and stable as required in Section 41, and is entitled to protection under the provisions of Section 42 of the Plant Variety Act.						
	Applicant(s) is (are) informed that false	representation herei	n can jeopardize prote	ection and result in p	enalties.		
	Jan. 5, 1982		Frank		. /		
	(DATE)		- I sank	IGNATURE OF APPLI	CANT	»."	
					•		

POPCORN

BIG RED

13A Exhibit A

Strawberry popcorn was the initial parent. It was planted adjacent to open pollenated yellow popcorn. Selected red grains from one yellow ear. Kernels were planted individually but allowed to cross pollinate. All red ear selections were planted and cross pollination allowed. One year used detassel method of male - female. Then corn was planted with no record male - female. In 1979 crop was all red with good size. Continued planting 1980 and 1981 and crop continued all red with nice size.

Now appears stable as to color.

POPCORN

BIG RED

13A Exhibit A

Strawberry popcorn was the initial parent. It was planted adjacent to open pollenated yellow popcorn. Selected red grains from one yellow ear. Kernels were planted individually but allowed to cross pollinate. All red ear selections were planted and cross pollination allowed. One year used detassel method of male - female. Then corn was planted with no record male - female. In 1979 crop was all red with good size. Continued planting 1980 and 1981 and crop continued all red with nice size. Corn has been stable both as to color and all varietal characters for these three generations.

There are variants that occur, these are smaller stalks with smaller cobs. They occur about 1 or 2 per 50 plants or be-

tween 2 to 4 per-cent.

13 B. Exhibit B.

Novelity Statement

Big Red is most similar to South American Giant Pop-corn.

Big Red is similar in size of grain to South American Giant yellow popcorn. While the yellow is approximately 2 feet shorter in height of stalk its leaf structure and color are similar to Big Red. It has a strong 2 ear tendency like Big Red but the ears are 3 to 4 inches longer. The yellow is superior in popping size also having less hull when popped. Their growing seasons are almost identical from sprout to maturity.

Big Red does however have a red pericap while the South American Giant Yellow has a yellow pericap.



EXHIBIT C

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, POULTRY, GRAIN & SEED DIVISION BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY

Frank Holland Address (Street and No. or R.F.D. No., City, State, and ZIP Code) 2615 Old Logan Rd, Rth B Lancaster, Onio 43130 Place the appropriate number that describes the varietal character of this variety in the boxes below. Place a zeto in first box (*-18- 0 8 9) or [0 9) when number is either 99 or less or 9 or less. 1. TYPE: 1. SWEET 2 - DENT 3 - FLINT 4 - FLOUR 5 - POP 6 - ORNAMENTAL 2. REGION WHERE BEST ADAPTED IN THE U.S.A: 7. SOUTHCENTRAL 6 - SOUTHCENTRAL 7 - MOST REGIONS 3. MATURITY (in Region of Best Adaptability): (Under "comments" (pg. 3) state how heat units were calculated.) DAYS FROM EMERGENCE TO BOX OF PLANTS IN SILK DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY 1. DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE 2. PLANT: R/5 * */3/82** 1. SOUTHCENTRAL 6 - SOUTHCENTRAL 7 - MOST REGIONS 3. MATURITY (in Region of Best Adaptability): (Under "comments" (pg. 3) state how heat units were calculated.) DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY 1. DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE 2. PLANT: R/5 * */3/82** 1. PLANT: Number of Tillers: Number of Tillers: Number of Tillers: 2. 1 - NONE 2 = 1-2 3 = 2-3 4 = >3 3. 1 - SINGLE 2 - SLIGHT TWO-EAR TENDENCY 3 - STRONG TWO-EAR TENDENCY 4 - THREE-EAR TENDENCY Cytoplesm Type: 1. PLANT 1 - NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER (Specify) 5. LEAF (Field Corn Inbred Exemples Gwen): Color:	CORN (ZEA MAYS)	
ADDRESS (SHAME AND No. CIP.). State, and 2IP Code 2C15 O14 LOGAN RR, Rt. (R. Rt.) (8 LANCASTER, COLIO . 43130 Place the appropriate number that describes the varietal character of this variety in the boxe below. Place a zero in first box (Fd. [0] 8] 5] or [0] 9) when number is either 99 or less or 9 or less. 1 rype: 5	NAME OF APPLICANT(S)	FOR OFFICIAL USE ONLY
2615 014 Logar Rd, Rth Lanobeter, Ohio 43130 Place the appropriate member that describes the varietal character of this variety in the boxes below. Place a zero in first box (**A** [0] 813) or [0] **) when number is either 99 or less or 9 or less.* 1. 179E: 1 - SWEET		. <u> </u>
Place the appropriate number that describes the varietal character of this variety in the boses below. Place a zero in first box (**-4** [0] 15**] or less of yor less of		
Place the appropriate number that describes the varietal Character of this variety in the boxes below. Place a zero in first box (*-as O B T) or O T) when number is either 57 or less or 9 or less. 1 TYPE: 1		
Place a Acto in first box (c-b 0 8 9 or 0 9 9 when number is cither 99 or less or 9 or less. Type:		Big Ked
1. TYPE:	Place the appropriate number that describes the varietal character of this variety in the	boxes below.
2. REGION WHERE BEST ADAPTED IN THE U.S.A.: 1 = NORTHWEST		9 or less.
2. REGION WHERE BEST ADAPTED IN THE U.S.A.: 1 - NORTHWEST 2 = NORTHKENTRAL 3 - NORTHEAST 7 - MOST REGIONS 3. MATURITY (In Region of Best Adaptability): (Under "comments" (pg. 3) state how heat surface where the state is were calculated by the state that the state of the state that were calculated by the stat	1. TYPE:	
1 = NORTHEAST 2 = NORTHEAST 7 = MOST RECIONS 3. MATURITY (in Region of Best Adaptability): (Under "comments" (pg. 3) state how heat units were calculated.) (A plant: (D) 9 4 5	5 1 = SWEET 2 = DENT 3 = FLINT 4 = FLOUR 5 = P	OP 6 = ORNAMENTAL
3. MATURITY (In Region of Best Adaptability): (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder "comments" (pg. 3) state how heat units were calculated.) (Inder units)	2. REGION WHERE BEST ADAPTED IN THE U.S.A.:	
### A J DAYS FROM EMERGENCE TO 50% OF PLANTS IN SILK T DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO OPTIMINE DIVINIS DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE D DAYS FROM 50% SILK TO HARV	171	4 = SOUTHEAST
4 3 DAYS FROM EMERGENCE TO 50% OF PLANTS IN SILK 7 2 DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY 1 2 9 0 HEAT UNITS 1 2 9 0 HEAT UNITS 1 2 9 0 HEAT UNITS 4. PLANT: 287 R/5 H/21/8 CM. HEIGHT (To tassel tip) 137 CM. LENGTH OF TOP EAR INTERNODE Number of Tillers: Number of Ears Per Stalk: 2 1 = NONE 2 = 1 - 2: 3 = 2 - 3 4 = > 3 3 1 ** SINGLE 2 - SLIGHT TWO-EAR TENDENCY 4 ** THREE-EAR TENDENCY 5 = OTHER (Specify) 5. LEAF (Field Corn Inbred Examples Given): Color: 3 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 - DARK GREEN (B14) 4 = VERY DARK GREEN (K166 Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = <30° 2 = 30 - 60° 3 = > 60° 2 1 = LIGHT (W22) 2 = MEDIUM (WF9) 3 - MARY (OH7L) 3 = MARY (OH7L) 3 = MARY (PA11) Length: Length: 1 1 CM. WIDEST POINT OF EAR NODE LEAF 1 1 NUMBER OF LEAVES PER MATURE PLANT	3. MATURITY (In Region of Best Adaptability): (Under "C	comments" (pg. 3) state how
7 2 DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY 1 2 9 0 HEAT UNITS	THE DAYS EDOMENTED TO THE TO THE STATE OF TH	
3 1 DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE 1 2 9 0 0 2 5 7 1 7 7 7 7 7 7 7 7		4 5
4. PLANT: 1. 5	7 2 DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY 1 2	9 0 HEAT UNITS
1 5 0 8 CM. HEIGHT (To tassel tip)	3 1 DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE 0 2	5 7 HEAT UNITS
1 0 8 CM. HEIGHT (To tassel tip) CM. LENGTH OF TOP EAR INTERNODE Number of Tillers: Number of Ears Per Stalk:		
Number of Tillars: Number of Ears Per Stalk: 2		
Number of Tillers: Number of Ears Per Stalk:	1 1 5 10 18 CM, HEIGHT (To tassel tip)	O 4 CM. EAR HEIGHT (TO BEST OF OST)
2 1 = NONE 2 = 1-2 3 = 2-3 4 = > 3 1 = SINGLE 2 = SLIGHT TWO-EAR TENDENCY 3 = STRONG TWO-EAR TENDENCY 4 = THREE-EAR TENDENCY Cytoplasm Type: 1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER (Specify) 5. LEAF (Field Corn Inbred Examples Given): Color: 3 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN (B14) 4 = VERY DARK GREEN (K166) Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = LIGHT (W22) 2 = MEDIUM (WF9) 3 = HEAVY (OH26) Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 3 = MANY (PA11) Width: Length: 1 1 CM. WIDEST POINT OF EAR NODE LEAF 1 1 1 CM. EAR NODE LEAF	1 0 CM. LENGTH OF TOP EAR INTERNODE	
5. LEAF (Field Corn Inbred Examples Given): Color: 3 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN (B14) 4 = VERY DARK GREEN (K166) Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = C 30° 2 = 30-60° 3 = > 60° 2 1 = LIGHT (W22) 2 = MEDIUM (WF9) 3 = HEAVY (OH26) Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 3 1 = ABSENT (OH51) 2 = FEW (OH56A) Width: Length: 1 1 CM. WIDEST POINT OF EAR NODE LEAF 1 1 NUMBER OF LEAVES PER MATURE PLANT	3=STRONG TWO	
5. LEAF (Field Corn Inbred Examples Given): Color: 3	1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER	(Specify)
Color: 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN (B14) 4 = VERY DARK GREEN (K166) Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 3 1 = ABSENT (OH51) 2 = FEW (OH56A) 3 = MANY (PA11) Width: Length: 1 1 CM. WIDEST POINT OF EAR NODE LEAF 1 1 1 CM, EAR NODE LEAF		
1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GREEN (B14) 4 = VERY DARK GREEN (K166) Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° 2 1 = LIGHT (W22) 2 = MEDIUM (WF9) 3 = HEAVY (OH26) Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 3 1 = ABSENT (OH51) 2 = FEW (OH56A) Width: Length: 1 1 CM. WIDEST POINT OF EAR NODE LEAF 1 1 CM. EAR NODE LEAF	5. LEAF (Field Corn Inbred Examples Given):	
Angle from Stalk (Upper half): Sheath Pubscence: 2 1 = < 30° 2 = 30-60° 3 = > 60° Calculate	Color:	•
2 1 = <30° 2 = 30-60° 3 = > 60° Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 3 1 = ABSENT (OH51) 2 = FEW (OH56A) 3 = MANY (PA11) Width: Length: 1 1 CM. WIDEST POINT OF EAR NODE LEAF 1 1 1 CM. EAR NODE LEAF	1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE	EN (B14) 4 = VERY DARK GREEN (K166
Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 3 1 = ABSENT (OH51) 2 = FEW (OH56A) Width: Length: 1 1 CM. WIDEST POINT OF EAR NODE LEAF 1 1 1 CM, EAR NODE LEAF	Angle from Stalk (Upper half): Sheath Pubscence:	
Marginal Waves: Longitudinal Creases: 2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 3 1 = ABSENT (OH51) 2 = FEW (OH56A) Width: Length: 1 1 CM. WIDEST POINT OF EAR NODE LEAF 1 1 NUMBER OF LEAVES PER MATURE PLANT		
Marginal Waves: Longitudinal Creases: 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 3 1 = ABSENT (OH51) 2 = FEW (OH56A) 3 = MANY (PA11) Width: Length: 1 1 CM. WIDEST POINT OF EAR NODE LEAF 1 1 NUMBER OF LEAVES PER MATURE PLANT		•
2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 3 1 = ABSENT (OH61) 2 = FEW (OH66A) 3 = MANY (PA11) Width: Length: 1 1 CM. WIDEST POINT OF EAR NODE LEAF 1 1 NUMBER OF LEAVES PER MATURE PLANT		, ,
Width: Length: CM. WIDEST POINT OF EAR NODE LEAF I 1 1 CM. EAR NODE LEAF I 1 1 NUMBER OF LEAVES PER MATURE PLANT		
Width: Length: CM. WIDEST POINT OF EAR NODE LEAF 1 1 1 CM. EAR NODE LEAF 1 1 1 CM. EAR NODE LEAF	161	
1 1 cm. WIDEST POINT OF EAR NODE LEAF 1 1 number of leaves per mature plant		
1 1 NUMBER OF LEAVES PER MATURE PLANT	<u>gara, me</u> sa selaka da kacamatan	
	1 1 cm. WIDEST POINT OF EAR NODE LEAF	R NODE LEAF
1 FORM LPGS-470-28 (3-79) (Formerly Form GR-470-28 (2-74) which may be used) Page Lof 3	1 1 NUMBER OF LEAVES PER MATURE PLANT	
Control of the state of the sta	FORM LPGS.470.28 (3.79) (Formerly Form CR.470.28 (2.74) which may be used)	Penal of 2

	Vik / inigat takts	The second second	নাম্যান ক্ষিত্ৰ কৈ জুলা মানকে মুকুজুলা ৰ	医化乳溶解 引起身 医生物结合物 化甲	PT (AM COMP)	 			3.8 38 S
6.	TASSEL:								٠
	08	11	子表 医水溶素 产品商 数十分		•				
	The state of the s		FLATERAL BRAN	the state of the s	The second secon	ri Si Cyri e ya kabina			-
11.7		7		18 M M2	ويستفير الأميدارة		* * * * *		
		le from Central	Spike:		Penduncle Length:	:		4	
	3	1 = < 309	2 = 30-40°	3 = > 45°	0 7	CM. FROM TOP	LEAF TO BAS	AL BRANCHE	ES
		THROMETHY.	1. 多十五百分分数3	As Series (Maries)		Vestua forest	1. 多量量。	7597795 ·	
	Pollen Shed	:			Service of	# 			
	2	\$30 m2			Complete (Property Commence)		,	•	
	L	1 = LIGHT (V	WF9)	2 = MEDIUM	, ,	/(KY21)			
		a te transfer		m in College and		Professional Control	그런 사람이 사람이 되었다. 	, federal in the control of the cont	
		Anther Color:)						
				W 2 = PINI	K golinik p o leM e E) 4=PU	HPLE	5 = GREEN	•
	1	Glume Color:) 6=OIHER	(Specify)			-		
	Pollen Resto	oration for Cyte	onlasms to = Not T	ested, 1 = Partial, 2 = G	nod)				
	- (-	•	53.00, 1 10.1.0., 2				•	
	[A] "T"		"s" 🔼	"c"	THER (Specify Cyto	nleem and degrees	of restoration)		
	י עטו	O	" D	٠ ـ ـ ـ ـ ـ ـ ـ ـ ـ ـ ـ ـ ـ ـ ـ ـ ـ ـ ـ	THEIR (Specify Cyte	Apiasiii and degrees	Of restoration,		
						•	•		
7.	. EAR (Husk	ced Ear Data Ex	cept When Stated (Otherwise):					
,			<u></u>		 				
1	20	CM LENGTH		M. MID-POINT	1 6 2	GM. WEIGHT	*		
		en en de la companya	ان حسی	WMEIEN		e de la companya de La companya de la co			
	Kernel Rows	5 :				art.			
1	2	1 - INDICTIN	ICT	- DISTINCT	20	NUMBER			
		1 - INDISTIN	.01	- DISTINCT		and the second			1.
					•	•	•		-
	2	1 = STRAIGH)T 2=SI	LIGHTLY CURVED	3 = SPIRAL	L			
:			·						
	Silk Color (E	exposed at Silki	ng Stage):			-			
1	4	1 = GREEN	2 = PINK	3 = SALMO	N 4=R	ED.	•		
1							1.	•	
	Husk Color:						•		
1	Trusk Color.								
.	1	FRESH	1 = LIGH	IT GREEN	2 = DARK GRE	EN	3 = PINK		
ļ		DRY	4 = RED	5 = P	URPLE	6 = BUFF		-	
ŧ	6		<i>Y</i>			3 30. 1			٠
. 1		on: (Harvest S			Husk Leaf:		-		
	2 1=SH	ORT (Ears Exp	osed) 2 = MEDIUM Beyond Ear Tip)	M (Barely Covering Ear)		= SHORT (< 8 CM	/	UM (8-15 CM	4)
ı.		RY LONG (>			3 لنگنا	= LONG (> 15 C	:M)		
5	Shank:			v v	Position at Dry Hus	k Stage:			
Γ	07		4					•	
Ļ		CM LONG	NO. OF I	NTERNODES	1 =	= UPRIGHT 2	2 = HORIZONT	AL 3 = PEI	NDEN
7	Taper:		•		Drying Time (Unhu	ekod Farl	:		
ſ						sico Laij.	*		
Į	1	1 = SLIGHT	2 = AVERAGE	3 = EXTREME	2 1-	SLOW 2	2 = AVERAGE	3 = FA	ST
	KERNEL (Dr	ied):							
į		ar Mid-Point):	*					* * *	
	Size (From E			•					
		MM LONG		i					
		MM LONG	06	MM. WIDE	0 5 MM, THI	ск			
[06	MM. WIDE	0 5 MM, THI		· · · · · · · · · · · · · · · · · · ·	₩.	-

		<u>, , , , , , , , , , , , , , , , , , , </u>	<u> </u>	<u> </u>	820	0076	
8. KERNEL	L (Dried) :		•		-		
7	5 = BRO	MM	6 = LIGHT REC			4 = BRONZE	
1	Aleurone Color: 1 = HOM	ozygous	2 = SEGREC	ATING (Describe)_			
1	· · · · · · · · · · · · · · · · · · ·		. ***	4 = BROWN	5	≃ BRONZE	6 = RI
1					4 = PINK-ORAN	GE 5 = WHITE	CAP.
Endosperi	m Type:			·			
		Color: 1 = COLORLESS	SH AMY L OSE STAR HER (Specify) [©]	СН			
18	GM. WEIGHT /100 SEEDS (Insized Sample)		· .			
9. coB:	MM. DIAMETER AT MID-PO	INT					
Strength:	1 = WEAK 2 = STRO	NG		1 = WHITE 2 =		4 = BROWN R (Specify)	
10. DISEASE	RESISTANCE (O = Not Teste	d, 1 = Susceptible, 2	? = Resistant):				
		0 so 0 co 0 m/ d guess 1s	OUTHERN LEAF DRN SMUT AIZE DWARF MO	BLIGHT DSAIC at to Ohio	O SMU	FERIAL WILT	
	CORNBORER ROOTWORM (Northern)	O EARWORM O ROOTWORM	f (Western)	رکا	EET L E	O APHID	
. VARIETIE	S MOST CLOSELY RESEMB	LING THAT SUBMI	TTED FOR THE	CHARACTERS GIV	/EN:		
CHARACT Maturity	ER	VARIETY	'			VARIETY	
Plant Type			1)		
Ear Type		J		Usage			
C E T S	J.S. Department Agriculture. Corn: Culture, Processing, Processing, R.A., G.W. Beadle, ar The Mutants of Maize. 1968. (tringfield, G.H. Maize Inbred I	ducts. 1970 Avi Pub id A.C. Fraser. A Sui Crop Science Society Lines of Ohio, Ohio	mmary of Linkag of America. Ma A.E.S. Bul. 831.	e Studies in Malze.Co dison, Wisconsin. 1959.	ernell A.E.S., Mem.	180. 1935.	•
		·		* .		 	rest
olumbus,	_						

Page 3 of 3

FORM LPGS-470-28 (3-79)

13D. Exhibit D

Additional description of Big Red

Stalk is taller and has more stockiness than the yellow. Tassel seems to tower above the stalk more than the yellow. Is a good number of 2 eared stalks. Basically the main difference is that the ears are red with good size.

Corn pops with a white color but has a red hull. Hull is more predominate than in yellow and taste is a little different.